## TRAUMATIC SPINAL CORD INJURY

Demographics at a Glance 2023




302,000

(Range from 255,000 to 383,000)

ESTIMATED PERSONS CURRENTLY LIVING
WITH SPINAL CORD INJURY

18,000

ESTIMATED NEW INJURIES EACH YEAR IN THE UNITED STATES





	56.1%	Non-Hispanic White
	24.7%	Non-Hispanic Black
	14.4%	Hispanic
	2.6%	Asian
•	1.7%	Other
	.5%	Native American

CAUSES OF INJURY
SINCE 2015



31 5%



**37.6%** Vehicular

31.5% Falls

15.4% Violence







8.3% Sports

3.6% Medical

3.6% Other

## NEUROLOGICAL LEVEL AND EXTENT OF INJURY

**SINCE 2015** 

47.1% Incomplete Tetraplegia

20.1%

Incomplete Paraplegia

19.8%

**Complete Paraplegia** 

12.5%

Complete Tetraplegia

.6%

Normal

© 2023 Board of Trustees, University of Alabama. This is a publication of the National Spinal Cord Injury Statistical Center offering content developed under a grant from the National Institute on Disability, Independent Living, and Rehabilitation Research (NIDILRR grant number 90SIMS0016). NIDILRR is a Center within the Administration for Community Living (ACL), Department of Health and Human Services (HHS). Content does not necessarily represent the policy of NIDILRR, ACL, HHS, and you should not assume endorsement by the Federal Government.

AVERAGE	TRENDS OVER TIME	1970s	<b>SINCE 2015</b>
	Age at Injury	29 Years Old	43 Years Old
	Acute Hospital Lengths of Stay	<b>24</b> Days	12 Days
K.	Rehabilitation Lengths of Stay	98 Days	31 Days